

New γ Doradus Stars from the Hipparcos Mission and Geneva Photometry

Laurent Eyer, Conny Aerts¹

*Instituut voor Sterrenkunde, Katholieke Universiteit Leuven,
 Celestijnenlaan 200 B, B-3001 Leuven, België*

Abstract. A search for new γ Dor stars was undertaken using the Hipparcos periodic variable star catalogue and the Geneva photometric database, leading to a list of 40 new candidates. We started a monitoring of the candidates which suited the observational window with the CORALIE spectrograph at the Swiss Euler Telescope for establishing a robust list of new γ Dor stars and studying line profile variations. We here present our long-term program.

1. Introduction

The γ Dor stars have amplitude variations up to 0.1 mag in Johnson V and periods ranging from 0.4 to 3 days (Kaye, these proceedings). We searched in two databases for finding new members of this class of variable stars. The first one is the Hipparcos main mission photometric database. It contains a mean of 110 measurements for 118 204 stars brighter than 12.4 and is magnitude complete up to 7.3-7.9 depending on the galactic latitude b . As the sampling is ruled by the scanning law of the satellite, it is not affected by the aliasing around 1/day, which might be a problem for detecting γ Dor stars.

The second scanned database is the Geneva photometric catalogue (Burki & Kienzie, these proceedings), it counts 48 000 stars and 345 000 measurements in a seven colour system. The content of the Geneva catalogue is the reunion of more than 200 scientific programmes, including namely the Bright Star Catalogue south of $\delta < +20$.

2. Hipparcos main mission

Thousands new variable stars were discovered by the Hipparcos satellite. During the analysis of the Hipparcos photometry, stars from the Periodic Catalogue having accurate parallaxes and colours were plotted in the HR diagram (Eyer 1998). A clump of stars just at the cool lower edge of the δ Scuti instability strip was present and gave rise to a list of 15 candidates (excluding redundant cases from the other studies). A clump is also present when plotting the variable stars of the Hipparcos Unsolved Catalogue.

¹Postdoctoral Fellow, Fund for Scientific Research, Flanders; e-mail: conny@ster.kuleuven.ac.be

3. Hipparcos main mission and Geneva photometry

A systematic search for finding new γ Dor stars was undertaken in the Hipparcos periodic variable star catalogue using also Geneva photometry and performing a multivariate discriminant analysis. This study led to a list of 14 new γ Dor stars (Aerts et al. 1998). This method is stricter since information on amplitude, period, physical parameters and multiperiodic behaviour were taken into account.

4. The search in Geneva photometry

Finally the Geneva photometric database was scanned to find F dwarf stars with high standard deviation. Eleven candidates were then measured with the 70-cm Swiss telescope, resulting in about 1000 photometric measurements, which are under study (Eyer & Aerts, in preparation). It turns out that about half suspected stars might be constant stars.

5. Spectroscopic measurements

In order to confirm the pulsational character of these stars, new spectra have been taken with the CORALIE spectrograph on the 1.2-m Swiss telescope at ESO-La Silla Observatory. The photometry and spectroscopy are necessary steps since we want to establish a robust list of new γ Dor stars. Up to now 22 stars have been measured, the strategy consist of taking at least five spectra of each candidate. Among the stars, some are binaries, some are fast rotators and some show clear line profile variations (cf. Fig. 1). Some stars are too faint for the telescope size, thus correlation techniques are used to lower the noise level. The following step is to accumulate photometry and spectra for promising candidates in order to perform mode identification.




Figure 1. Line profile variations of the star HD 14940.

References

- Aerts, C., Eyer, L., & Kestens, E. 1998, A&A337, 790
 Eyer, L. 1998, PhD Thesis, Geneva University